

**American Journal on Intellectual and Developmental Disabilities**  
**Chronic Health Conditions among Adults with I/DD in a State Medicaid System**  
--Manuscript Draft--

|                                     |   |
|-------------------------------------|---|
| <b>Manuscript Number:</b>           | AJIDD-D-23-00044R3  |
| <b>Article Type:</b>                | Research Report   |
| <b>Keywords:</b>                    | autism spectrum disorder, intellectual and developmental disability, physical health, Medicaid, health disparities  |
| <b>Corresponding Author:</b>        | Kiley McLean, MSW, MEd<br>Drexel University AJ Drexel Autism Institute<br>Madison, Wisconsin UNITED STATES  |
| <b>First Author:</b>                | Kiley McLean, PhD, MSW, MEd   |
| <b>Order of Authors:</b>            | Kiley McLean, PhD, MSW, MEd<br>Lauren Bishop, PhD, MSW  |
| <b>Manuscript Region of Origin:</b> | UNITED STATES   |
| <b>Abstract:</b>                    | Despite a growing number of adults with intellectual and developmental disabilities (I/DD) and documented risk for adverse outcomes as they age, little is known about the health and healthcare patterns of adults with different I/DD throughout adulthood. This study uses Wisconsin Medicaid claims data to characterize health conditions among adults with I/DD. Results indicate high prevalence of asthma, diabetes, heart disease, and hypertension. Heart disease rates were particularly high, having been observed among 39% of autistic adults, 64% of autistic adults with intellectual disability, 67% of adults with Down syndrome, and 75% of adults with intellectual disability only. Given there are no known biological differences underlying increased morbidities among most people with I/DD, developing inclusive prevention measures should be prioritized in future research. |

### Abstract

Despite a growing number of adults with intellectual and developmental disabilities (I/DD) and documented risk for adverse outcomes as they age, little is known about the health and healthcare patterns of adults with different I/DD throughout adulthood. This study uses Wisconsin Medicaid claims data to characterize health conditions among adults with I/DD. Results indicate high prevalence of asthma, diabetes, heart disease, and hypertension. **Heart disease rates were particularly high, having been observed among 39% of autistic adults, 64% of autistic adults with intellectual disability, 67% of adults with Down syndrome, and 75% of adults with intellectual disability only.** Given there are no known biological differences underlying increased morbidities among most people with I/DD, developing inclusive prevention measures should be prioritized in future research.

*Keywords:* autism spectrum disorder, intellectual and developmental disability, physical health, Medicaid, health disparities

### Chronic Health Conditions among Adults with I/DD in a State Medicaid System

Medicaid provides health care coverage to more than 40 million people in the United States (Bruen & Holahan, 2001). Medicaid serves low-income adults, children, pregnant women, elderly adults, and disabled individuals\* (Centers for Medicare & Medicaid Services, 2019). According to the Medicaid and CHIP Payment and Access Commission (MACPAC) 2018 report to Congress, over 10 million people qualify for Medicaid on the basis of disability. This population includes disabled adults that present at birth and those who have acquired disabling conditions through illness, injury, or trauma (Centers for Medicare & Medicaid Services, 2019).

#### *Wisconsin's Medicaid Program*

Medicaid and the coverage and services provided to its beneficiaries vary from state to state. The way in which a state structures and implements its Medicaid program is greatly influenced by the program's unique funding structure. Medicaid is funded jointly by states and the federal government (Snyder & Rudowitz, 2015). This state and federal partnership is governed by the federal medical assistance percentage (FMAP), which guarantees federal match funds to states for qualifying Medicaid expenditures (e.g., payments states make for covered Medicaid services provided by qualified providers to eligible Medicaid beneficiaries) (Snyder & Rudowitz, 2015). Thus, FMAP, and its corresponding state Medicaid program, varies from one state to the next. Wisconsin's FMAP is set at 66.30% for fiscal year 2023 (KFF, n.d.). This standard FMAP formula applies to the vast majority of Medicaid spending, though there are exceptions that provide higher match rates, many of which were enacted through the Affordable Care Act of 2010 (ACA) (Snyder & Rudowitz, 2015). The ACA broadened Medicaid's role, providing healthcare coverage to many adults previously uninsured and increasing the state's

---

\* The authors of this manuscript chose to use identity-first language (i.e., disabled person, autistic person) to respect the preferences of many of those within the disability community.

FMAP (Snyder & Rudowitz, 2015). That said, the Supreme Court ruled the ACA's Medicaid expansion optional for states and as of December 2022, eleven states have not adopted it (KFF, 2023). Wisconsin is one of these eleven.

Wisconsin has not implemented the ACA's Medicaid expansion, choosing to limit Medicaid eligibility and forgo the enhanced FMAP for their expansion populations (KFF, 2023). States that have adopted the ACA's Medicaid expansion provide coverage to nearly all adults with incomes up to 138% of the Federal Poverty Level (KFF, 2023). In contrast, Wisconsin covers adults up to 100% of the Federal Poverty Level (KFF, 2023). Individuals in Wisconsin must meet these financial requirements set by the state in addition to categorical eligibility requirements in order to be eligible for its Medicaid program. As prior mentioned, many meet the requirements for categorical eligibility on the basis of disability. National estimates from 2008-2016 indicated that 81.7% of autistic adults who were eligible for Medicaid were through disability compared to 8.8% through poverty (Roux et al., 2023).

Nationally, Medicaid beneficiaries enrolled through disability pathways include those with physical disabilities, intellectual and developmental disabilities (I/DD), and serious behavior disorders and mental illness (MACPAC, 2018). Among these groups, individuals with I/DD are a primary group of beneficiaries. An individual with an I/DD, such as autism spectrum disorder, Down syndrome, and intellectual disability, can qualify for Medicaid in Wisconsin through three pathways: 1) they receive Supplemental Security Income (SSI) benefits; (2) they purchase Medicaid health benefits through the Medicaid Purchase Plan; or (3) they have income at or below 100% of the Federal Poverty Level (KFF, n.d.). In summation, an understanding of the complexity and variability of state Medicaid programs like Wisconsin's and their I/DD systems provides essential context for our interpretation of healthcare research. Indeed, the

national landscape of Medicaid programs is complex. While efforts have been made to improve outcomes of disabled individuals on the federal level, it is important to understand how specific states are contributing to or addressing health disparities in adults with I/DD. Policies with meaningful improvements for instance may then be implemented in similar settings. Therefore, an investigation into a state Medicaid program like Wisconsin's may assist in reshaping future healthcare policy, including in potential adoption of the ACA expansion in states that have not done so already.

#### *Medicaid Research with Adults with I/DD*

Thus far, Medicaid research in relation to individuals with I/DD has been limited, and primarily focuses on children and those with autism spectrum disorder (Mandell et al., 2010; Wang & Leslie, 2010). Recent research has examined the changing Medicaid enrollment patterns in autistic adult Medicaid beneficiaries and distinguishes this community as a primary and growing group of Medicaid beneficiaries. Some states have found a consistent influx of new autistic young adults into the state Medicaid system over the past decade, likely corresponding to the increased identification of autistic children (Rubenstein & Bishop, 2019).

Research has also suggested that there may be county-level variation in the prevalence of Medicaid-enrolled autistic children (Mandell et al., 2010). Medicaid-enrolled autism spectrum disorder prevalence varies greatly as a function of several county characteristics. Counties with lower per-student education expenditures, more students, a greater proportion of students in special education, higher per capita number of pediatricians and pediatric specialists, and a greater proportion of Medicaid enrollees and White residents had higher Medicaid prevalence (Mandell et al., 2010). Therefore, it seems that not only do states differ in their implementation of Medicaid policies, but counties within these states may differ as well. It's been hypothesized

that these differences in county-level Medicaid implementation may relate to availability of educational resources, counties taking advantage of their state's Medicaid policies, and variation in access in rural and urban areas (Mandell et al., 2010).

*Chronic Health Conditions in Adults with I/DD*

Adults with I/DD may be at increased risk for chronic health conditions as they age. In its 2010 report on the status of the challenges presented by chronic health conditions, the World Health Organization noted that non-communicable conditions, such as cardiovascular disease, diabetes, cancers, and chronic respiratory diseases, accounted for nearly two-thirds of all deaths worldwide (WHO, 2011). In the United States, chronic health conditions are the main causes of poor health and early mortality, and they account for the highest proportion of healthcare expenditures (Bauer et al., 2014). Research suggests that these chronic health conditions are a result of certain risk factors including tobacco use, poor diet, physical activity, excessive alcohol consumption, uncontrolled high blood pressure, and hyperlipidemia (Bauer et al., 2014; Fine et al., 2004). Of these risk factors, adults with I/DD are associated with high rates of physical inactivity, obesity, and smoking (Havercamp & Scott, 2015; Ranjan et al., 2018). Autistic children and adolescents have been found to have higher rates of obesity or being overweight than non-autistic youth, which could increase the risk of conditions such as diabetes, cardiovascular disease, and cancer (Ranjan et al., 2018). Perhaps consequently, a recent study using Medicaid data to compare health outcomes among middle-aged and older autistic adults (48-88 years old) with and without co-occurring intellectual disability found high prevalence of most health conditions including immune conditions, cardiovascular disease, sleep disorders, gastrointestinal disorders, neurologic conditions, and psychiatric disorders (Bishop-Fitzpatrick & Rubenstein, 2019). A study of over 1500 autistic adults, mean age 29, found elevated rates of

diabetes, hypertension, and hyperlipidemia, but marginally lower rates of cancer, compared to non-autistic adults in their sample (Croen et al., 2015). It should be noted that decreased risk of certain health conditions, like cancer, have been documented in some studies with autistic adults and other developmental disabilities (Bishop-Fitzpatrick et al., 2018). This could be attributed to disparities in screening and diagnosis, or potential protective factors among this group (Elder et al., 2016). Further research will be needed in this area.

Fortunately, many of the risk factors for chronic disease prevalent among adults with I/DD are nongenetic and potentially modifiable or preventable (Fine et al., 2004). It should be noted, however, that adults with Down syndrome do have documented biological differences compared to other disabled and non-disabled adults (Landes et al., 2020). Those with Down syndrome, or Trisomy 21, have a full or partial extra copy of chromosome 21 (Landes et al., 2020). This additional genetic material can alter the course of development and cause characteristics associated with Down syndrome in adults. For example, many people with Down syndrome experience craniofacial alterations and muscle hypotonia, developmental delays, cognitive impairment, and increased prevalence of some medical conditions including congenital heart disease and early onset Alzheimer's disease (Asim et al., 2015).

Although the underlying genetic cause of Down syndrome is the same, resulting pathologies and their severity vary from person to person (Lana-Elola et al., 2011). Due to advances in diagnosis and treatment of adults with Down syndrome for example, the overall health and mortality disadvantage for this group has considerably improved in the 20th century (Landes et al., 2020). A growing body of research has suggested that contrary to the generally accepted belief, cognitive decline by middle age for adults with Down syndrome, including risk for developing Alzheimer's disease, is not inevitable and can be modified by a number of factors

(Zigman et al., 2008). Societal factors including deinstitutionalization and medical factors including improved nutritional and public healthcare practices have resulted in these extensions of life expectancy and improvements in overall health in adults with Down syndrome (Zigman et al., 2008).

As such, differences in physical health conditions among adults with I/DD, including those with Down syndrome, may be evidence of health disparities (i.e., health problems that are unjust and avoidable) and perhaps can explain the two- to three- decades decreased life expectancy estimated in adults with I/DD (Piven & Ravins, 2011; Bishop-Fitzpatrick & Rubenstein, 2019). The preliminary findings from these studies lay groundwork for research into health disparities among adults with I/DD.

Ultimately, very little is known about the unique health and healthcare patterns of individuals with I/DD throughout their adulthoods and the ways in which different state Medicaid programs address or contribute to these disparities (Piven & Rabins, 2011; Anderson et al., 2013). Most studies to date that examine chronic health conditions of adults with I/DD compare autistic children and young adults to those without disabilities and find greater prevalence of most health conditions (Croen et al., 2015; Weir et al., 2022). It is unclear, however, how health conditions develop across age groups and how they compare across those with different I/DD. Therefore, it is critical to use existing systems-level data to characterize the health status of this population. More specifically, assessing the prevalence of chronic health conditions in adults with different I/DD can help to inform prevention efforts, anticipate and prepare for service use, lessen healthcare expenditures, and ultimately, reduce health disparities.

### *Purpose*

Given the urgency for research in this area, one of the best tools for efficiently investigating health in an entire health system, without requiring years of follow-up, is the use of Medicaid claims data. Medicaid claims have been successfully used in research related to autistic children, adolescents, and young adults (Mandell et al, 2010; Wang & Leslie, 2010). And more recently, has been used to characterize health outcomes in middle-aged and older autistic adults (Vohra et al., 2017; Schott et al., 2022; Bishop-Fitzpatrick & Rubenstein, 2019). To our knowledge, no studies to date have used Medicaid claims data to characterize and compare health outcomes among young, middle-aged, and older adults with different types of I/DD. Thus, this study presents a novel approach to identifying and characterizing health disparities among adults with I/DD across the remainder of their lifespans.

For the purposes of this investigation, four of the most common chronic diseases in older adults that could be measured with our existing Medicaid data were chosen: asthma, diabetes, heart disease, and hypertension. Existing research suggests that autistic adults are at significant risk for developing these conditions in mid to later life (Tyler et al., 2011; Croen et al., 2015). How these progress with age, as well as compare to adults with other I/DD, needs further investigation.

The overall purpose of this study is to use state-level Medicaid claims data to characterize health outcomes among Medicaid beneficiaries, aged 21+, with I/DD. More specifically, we used Wisconsin Medicaid claims from 2008-2018 to examine the prevalence of chronic health conditions (asthma, diabetes, heart disease, and hypertension) among adults with autism only, autism and intellectual disability, Down syndrome only, and intellectual disability only, controlling for age, gender, race and ethnicity, and rural or urban county designation. We hypothesized that all groups would have high rates of each physical health condition and that

rates would rise as adults aged. Further, we hypothesized that those with intellectual disabilities would have greater prevalence of all conditions.

## Methods

### *Data Source*

We obtained de-identified Wisconsin Medicaid claims data from January 1, 2008, to December 31, 2018, through a limited data use agreement with the Wisconsin Department of Health Services. Adults (21+) were included in the data provided if they had two Medicaid claims for an I/DD on two different days during their entire lifetime period of Medicaid enrollment. Demographic information (age, race/ethnicity, gender, county of residence), medical claims (dental, home health, long term care, pharmacy and professional, inpatient, outpatient, and crossover claims), and the paid amount for each medical claim were included in the data. Corresponding ICD-9 and ICD-10 codes were provided for each claim, as well as Clinical Classification Software (CCS) codes for all physical and psychiatric health conditions.

### *Sample*

Adults (21+) were included in these data if they had two Medicaid claims for an I/DD on two different days during their entire lifetime period of Medicaid enrollment. Participants with at least 2 ICD-9 or ICD-10 codes for autism spectrum disorder (299.0, 299.00, 299.01, 299.8, 299.80, 299.81, 299.9, 299.90, 299.91 or F84.0, F84.5, F849), intellectual disability (317, 318, 318.0, 318.1, 318.2, 319 or F70, F71, F72, F73, F78, F79), and/or Down syndrome (758.0 or Q90, Q90.1, Q90.2, Q90.9) comprised the sample. Prior research has suggested that using two claims rather than one significantly increases the specificity of conditions studied and helps to ensure that the claim itself is not a rule out for another diagnosis (i.e., the diagnosis is recorded for a suspected condition but is not the primary or final diagnosis; Rector et al., 2004). This

method of sampling is also consistent with past work examining I/DD in claims and other electronic health records data in adults (Bishop-Fitzpatrick & Rubenstein, 2019; Vohra et al., 2017).

We classified Medicaid beneficiaries in these data into four separate diagnostic groups: (1) autism spectrum disorder without intellectual disability; (2) autism spectrum disorder with intellectual disability; (3) Down syndrome; and (4) intellectual disability only. Most studies to date either group all adults with I/DD together in their analyses or compare autistic adults to non-autistic adults (Anderson et al., 2013). Only recently has research investigated prevalence estimates of certain health conditions stratified by diagnosis (autism, Down syndrome, and intellectual disability) (Ptomey et al., 2020). Given the heterogeneity of I/DDs, it is imperative to assess outcomes by I/DD type. This study seeks to add to this literature, recognizing that each type of I/DD present differently and that to improve the overall health of adults with I/DD, patterns and differences across diagnoses must be better understood.

Our original data included 24,149,448 total medical claims and 49,309 Medicaid beneficiaries with I/DD. Of these Medicaid beneficiaries with I/DD, 15,257 had medical claims during the study period (January 1, 2008-December 31, 2018). Thus, only those with two claims for I/DD and with medical claims during the study period were included in analysis (i.e., complete case analysis). Our final analytic sample includes 2828 adults with autism only, 1620 adults with autism and intellectual disability, 2040 adults with Down syndrome, and 8769 adults with intellectual disability only.

### *Chronic Health Conditions*

We examined all medical claims for a select group of health condition codes (asthma, diabetes, heart disease, hypertension) between January 1, 2008 to December 31, 2018. Each

ICD-9/ICD-10 code corresponds to a CCS diagnosis. The Agency for Healthcare Research and Quality H-CUP Clinical Classification Software (CCS) is a diagnosis and procedure categorization scheme used in projects analyzing data on diagnoses, including those tracking healthcare utilization comorbidity prevalence in autistic adults (Vohra et al., 2017). The CCS collapses ICD codes into a smaller number of clinically meaningful categories that are often more useful for presenting descriptive statistics on a population, rather than individual ICD codes. The CCS allows similar ICD codes to be grouped into an overarching diagnostic category for analysis. For example, the CCS category of hypertension includes essential hypertension, hypertension complication, and hypertension with complications and secondary hypertension. Thus, a CCS category may contain ICD codes with differing levels of severity. For the purposes of this study, we classified appropriate ICD-9 and ICD-10 codes into single-level CCS codes. Then, we grouped these single-level CCS codes into the multi-level CCS diagnostic categories for asthma, diabetes, heart disease, and hypertension. The CCS categories and their corresponding CCS and Medicaid codes can be found in Appendix B.

#### *Analytic Plan*

We first examined demographic variables provided in the enrollment data by I/DD status. Race/ethnicity were coded according to the condensed Medicaid categories for race and ethnicity (MACPAC, 2022). We summarized categorical variables by using frequencies and percentages and continuous variables through means and standard deviations. Based on the Wisconsin Office of Rural Health rural classifications, we coded all counties into rural or urban categories (Wisconsin Office of Rural Health, 2022). We estimated proportions of physical health conditions by diagnostic group and age category. We then graphed these proportions to highlight the differences in physical health condition claims for each diagnostic group across adulthood

(21+). Finally, given the high rates of physical health conditions in adults with intellectual disability in our sample, we conducted a post-hoc analysis to examine unadjusted and adjusted odds ratios of each physical health condition comparing those with intellectual disability to those without intellectual disability.

### Results

Demographic comparisons are presented in Table 1. Mean ages as well as age category frequencies are provided. Individuals within the intellectual disability only group made up 57.48% of the study population as well as had the highest proportion of older adults (50-64 and 65+). In comparison, 71.75% of individuals within the autism only group fell within the 21-34 age group. Of note, the number of autistic adults with and without intellectual disability diminished as age rose. For example, our sample had 2,029 autistic adults aged 21-34 and only 68 aged 65+. Similarly, our sample had 653 autistic adults with intellectual disability aged 21-34 and 148 aged 65+. Gender differences were most notable within the autism only group with 72.69% male. All groups had the highest proportion of non-Hispanic white adults compared to other racial and ethnic identities. Further, 69-75% of all groups had beneficiaries residing in urban counties.

[Insert Table 1 here.]

#### *Physical Health Conditions*

Tables 2a-d present the proportions of adult Medicaid beneficiaries with medical claims for asthma, diabetes, heart disease, and hypertension across diagnostic groups and age categories. In this sample, the prevalence of physical health conditions was high. Within the autism only group, 12.34% had claims for asthma, 18.81% for diabetes, 39.00% for heart disease, and 15.70% for hypertension. 72.25% of autistic adults aged 50-64 and 82.35% of autistic adults 65+

had claims for heart disease. Over half of those aged 50-64 had claims for diabetes as well.

Notably, prevalence of diabetes (30.31%), heart disease (63.73%), and hypertension (27.78%) were higher for those with autism and intellectual disability compared to those with autism only.

Prevalence rates for asthma were similar for those with Down syndrome only (12.70%) and intellectual disability only (13.35%). However, overall rates of heart disease and hypertension were statistically higher among these groups compared to those within the autism only or autism and intellectual disability groups. 66.91% of those within the Down syndrome only group and 75.29% of those within the intellectual disability only group had claims for heart disease. Across both diagnostic groups, over 80% of those aged 50-64 and 65+ had claims for heart disease. Further, 40.53% of those within the intellectual disability only group had claims for hypertension including over 50% of those aged 50-64 and 65+. Plots of the proportions of physical health conditions by age and diagnostic group can also be found in Figures 1a-d. The figures illustrate the overall prevalence of each health condition within each diagnostic group as well as the differences in prevalence across age groups.

[Insert Tables 2a-d here.]

[Insert Figures 1a-d here.]

In our post-hoc analyses comparing unadjusted and adjusted odds ratios of each of these physical health conditions (Tables 3a-b), we found statistically significant greater odds of heart disease and hypertension among all groups compared to those with autism only. Those within the autism and intellectual disability group had significantly ( $p < .001$ ) increased odds of diabetes (aOR=1.43, 95% CI [1.2, 1.7]), heart disease (aOR=1.76, 95% CI [1.5, 2.0]), and hypertension (aOR=1.41, 95% CI [1.2, 1.6]). Similarly, those with intellectual disability only had significantly ( $p < .001$ ) increased odds of diabetes (aOR=1.72, 95% CI [1.5, 1.9]), heart disease (aOR=2.12,

95% CI [1.9, 2.3]), and hypertension (aOR=1.83, 95% CI [1.6, 2.1]) as well. Finally, those with Down syndrome had significantly ( $p<.001$ ) increased odds of heart disease (aOR=1.68, 95% CI [1.5, 1.9]). Across all physical health conditions, those within the intellectual disability only group had the greatest odds of having a claim, compared to those with autism only. The adjusted odds ratios are graphed in Figure 2.

[Insert Tables 3a-b here.]

[Insert Figure 2 here.]

### Discussion

We used Wisconsin Medicaid claims data from 2008-2018 to examine the prevalence of specified chronic health conditions among adults with I/DD aged 21 and over. A small number of previous studies have used Medicaid claims data to identify a higher prevalence of physical health conditions among autistic adults compared to the general population and have suggested these health disparities will likely increase with age (Bishop-Fitzpatrick et al., 2018; Schott et al., 2022; Vohra et al., 2017). An even smaller number of studies have compared co-morbidity prevalence among middle-age and older autistic adults with and without intellectual disability through Medicaid claims and found heightened prevalence of physical health conditions in middle age and beyond, regardless of intellectual disability prevalence (Bishop-Fitzpatrick & Rubenstein, 2019). To our knowledge, no previous research has investigated the prevalence of health conditions among adults with I/DD in younger, middle, and older adults in a state Medicaid system.

First, our demographic comparisons point to interesting differences among adults with different I/DD diagnoses. Medicaid beneficiaries with autism only for instance were younger than other diagnostic groups, with over 70% falling within the 21-34 age group. Autism has a

relatively brief history as an identifiable and diagnosable condition compared to other I/DD; thus, this finding may represent the influx of new autistic young adults into our state Medicaid systems (Rubenstein & Bishop, 2019). It will be critical to track the changing enrollment patterns of autistic adults over the next few decades as larger cohorts of autistic adults are likely to age within the Medicaid system alongside the continually increasing identification of autistic children. Autistic adults with and without intellectual disability in our sample were also predominantly male (69-73%) whereas adults with Down syndrome only or intellectual disability only had a more balanced gender distribution. Interestingly, across all diagnostic groups approximately 70% of beneficiaries resided in urban Wisconsin counties. This may suggest that families with those with disabilities predominantly live in urban areas. However, rural communities may also face significant barriers regarding adequate availability of diagnostic, treatment, and support services for autistic individuals and those with other I/DD; thus, these data may be missing misdiagnosed or underdiagnosed adults in rural communities (Antezana et al., 2017).

Building on prior research in this area, our study found a high prevalence of physical health conditions among our adult Medicaid beneficiaries with I/DD. Compared to the other conditions, heart disease rates were particularly high across all groups including approximately 39% of autistic adults, 64% of autistic adults with intellectual disability, 67% of adults with Down syndrome, and 75% of adults with intellectual disability only. Though overall prevalence of heart disease in autistic adults was 39%, roughly 72% of autistic adults aged 50-64 and 82% 65+ had claims for heart disease during our study period, suggesting that health disparities among adults with I/DD are likely to increase with age. Considering these increases in prevalence alongside the drop in number of autistic adults from each rising age category to the

next, might point to mechanisms underlying early mortality in this population, though further research will be needed to parse out these causal relationships. Illustrations of the increase in health disparities with age can be found in Figures 3a-d. **Our other findings indicate that over 50% of autistic adults 65+ with and without intellectual disability had claims for hypertension.** Approximately 27% of adults within the Down syndrome only group and 40% of adults within intellectual disability only group had claims for diabetes during the study period. Approximately 11-13% of all adults in our study had claims for asthma during the study period.

In a review of the prevalence of chronic diseases among all adult Medicaid beneficiaries, Chapel and colleagues found that among 29 studies, prevalence estimates for enrollees aged 18-64 years were 8.8%-11.8% for heart disease, 17.2-27.4% for hypertension, 7.8%-19.3% for asthma, and 7.5%-12.7% for diabetes (2017). **Though prevalence of chronic diseases among all Medicaid beneficiaries in Wisconsin is unavailable, according to the National Center for Health Statistics, the death rate for heart disease in Wisconsin was 5% lower than the national rate (CDC, 2018).** These findings further highlight the heightened prevalence of chronic conditions among adults with different I/DD. For instance, heart disease rates in adults with intellectual disability are almost seven times greater than the estimates among all adult Medicaid beneficiaries overall.

Finally, our post-hoc analyses found that in controlling for age, gender, race and ethnicity, and urban or rural county designation, autistic adults with intellectual disability had significantly greater odds of having a claim for diabetes, heart disease, or hypertension compared to autistic adults without an intellectual disability. Compared to autistic adults without an intellectual disability, adults within the intellectual disability only group had the greatest odds of having any asthma, diabetes, heart disease, and hypertension claims. All other diagnostic groups

had significantly increased odds of having a claim for heart disease compared to autistic adults without intellectual disability.

### *Limitations*

These findings should be interpreted in light of several limitations. First, our sample represents Wisconsin Medicaid beneficiaries with a recorded I/DD from 2008 to 2018. Given changing diagnostic practices and improved availability of interventions, it is likely that many more adults with I/DD have entered this state Medicaid system since the end of the study period. Further, diagnoses of I/DD in Medicaid claims are not validated by a standardized clinical examination outside of examinations performed by the clinician at the time of entering the claim, possibly resulting in under and mis-diagnoses. Though measurement validity of adult I/DD diagnoses in Medicaid claims has yet to be determined, prior research has found 97-98% of autistic children with claims for autism did meet research criteria for autism (Fombonne et al., 2004). Further, because of missing data on race, this study could not closely draw comparisons among racial groups. Finally, the variability of state Medicaid programs described above may impact the generalizability of our findings. The low-income disabled population in Wisconsin may differ from other states or the overall population in the United States in some respects. Wisconsin-specific policies may also impact adult outcomes.

### *Implications*

Despite these limitations, this is among the first United States studies to examine patterns of chronic health conditions among adults with different types of I/DD in a state Medicaid program. Our analyses identify high prevalence of asthma, diabetes, heart disease, and hypertension among adults on Medicaid with autism, intellectual disability, and Down syndrome. Heart disease, in particular, may be a top area of concern for adults with I/DD, especially among

those with Down syndrome and other intellectual disabilities. Currently, there are no known biological differences underlying these increased morbidities prevalent among most people with I/DD (Al Dera, 2022; Mandell, 2018). For example, research has suggested that autism and its associated common comorbidities are of multi-origin and caused by a complex interaction of genetic, nutritional, and environmental factors (Al Dera, 2022; Tye et al., 2019). Adults with Down syndrome do have an underlying genetic cause; however, research suggests it does not fully account for their high prevalence of health disparities. With this context and with knowledge of current prevention measures for conditions like asthma, diabetes, heart disease, and hypertension, adverse health outcomes among adults with I/DD may be avoidable. It is therefore critical that future research investigate the causal mechanisms underlying these health disparities, and target interventions at both the individual and systems-level. Research should ensure prevention efforts for chronic health conditions are inclusive and evidence-based within I/DD communities. Inclusive, systems-level interventions built in partnership with all I/DD stakeholders have the potential to reduce healthcare costs and improve the overall health, quality of life, and longevity of adults with I/DD.

### References

- Al Dera, H. (2022). Cellular and molecular mechanisms underlying autism spectrum disorders and associated comorbidities: a pathophysiological review. *Biomedicine & Pharmacotherapy, 148*, 112688.
- Anderson, L. L., Humphries, K., McDermott, S., Marks, B., Sisirak, J., & Larson, S. (2013). The state of the science of health and wellness for adults with intellectual and developmental disabilities. *Intellectual and Developmental Disabilities, 51*(5), 385-398.
- Antezana, L., Scarpa, A., Valdespino, A., Albright, J., & Richey, J. A. (2017). Rural trends in diagnosis and services for autism spectrum disorder. *Frontiers in Psychology, 8*, 590.
- Asim, A., Kumar, A., Muthuswamy, S., Jain, S., & Agarwal, S. (2015). Down syndrome: An insight of the disease. *Journal of Biomedical Science, 22*(1), 1-9.
- Availability of race and ethnicity data for Medicaid beneficiaries* . MACPAC. (2022). Retrieved February 28, 2023, from [https://www.macpac.gov/wpcontent/uploads/2022/03/MACPAC-brief\\_Race-and-Ethnicity-Data-Availability.pdf](https://www.macpac.gov/wpcontent/uploads/2022/03/MACPAC-brief_Race-and-Ethnicity-Data-Availability.pdf)
- Bauer, U. E., Briss, P. A., Goodman, R. A., & Bowman, B. A. (2014). Prevention of chronic disease in the 21st century: elimination of the leading preventable causes of premature death and disability in the USA. *The Lancet, 384*(9937), 45-52.
- Bishop- Fitzpatrick, L., Movaghar, A., Greenberg, J. S., Page, D., DaWalt, L. S., Brilliant, M. H., & Mailick, M. R. (2018). Using machine learning to identify patterns of lifetime health problems in decedents with autism spectrum disorder. *Autism Research, 11*(8), 1120-1128.

Bishop-Fitzpatrick, L., & Rubenstein, E. (2019). The physical and mental health of middle aged and older adults on the autism spectrum and the impact of intellectual disability. *Research in Autism Spectrum Disorders*, 63, 34-41.

Bruen, B., & Holahan, J. (2001). Medicaid spending growth remained modest in 1998, but likely headed upward. *Kaiser Commission on Medicaid and the Uninsured Issue Paper*.  
<http://www.kff.org/medicaid/loader.cfm?url=/commonspot/security/getfile.cfm&pageId=13745>

Centers for Disease Control and Prevention. (2018, April 4). *Stats of the State of Wisconsin*.  
Centers for Disease Control and Prevention.  
<https://www.cdc.gov/nchs/pressroom/states/wisconsin/wisconsin.htm>

Centers for Medicare & Medicaid Services. (2019). *Medicaid state waiver program demonstration projects: Medicaid waivers and demonstrations list*.  
<http://www.medicare.gov/Medicaid-CHIP-Program-Information/By-Topics/Waivers/Waivers.html>

Chapel, J. M., Ritchey, M. D., Zhang, D., & Wang, G. (2017). Prevalence and medical costs of chronic diseases among adult Medicaid beneficiaries. *American Journal of Preventive Medicine*, 53(6), S143-S154.

Croen, L. A., Zerbo, O., Qian, Y., Massolo, M. L., Rich, S., Sidney, S., & Kripke, C. (2015). The health status of adults on the autism spectrum. *Autism*, 19(7), 814-823.

Elder, J. H., Brasher, S., & Alexander, B. (2016). Identifying the barriers to early diagnosis and treatment in underserved individuals with autism spectrum disorders (ASD) and their families: A qualitative study. *Issues in Mental Health Nursing*, 37(6), 412-420.

- Fine, L. J., Philogene, G. S., Gramling, R., Coups, E. J., & Sinha, S. (2004). Prevalence of multiple chronic disease risk factors: 2001 National Health Interview Survey. *American Journal of Preventive Medicine*, 27(2), 18-24.
- Fombonne, E., Heavey, L., Smeeth, L., Rodrigues, L. C., Cook, C., Smith, P. G., ... & Hall, A. J. (2004). Validation of the diagnosis of autism in general practitioner records. *BMC Public Health*, 4(1), 1-9.
- Havercamp, S. M., & Scott, H. M. (2015). National health surveillance of adults with disabilities, adults with intellectual and developmental disabilities, and adults with no disabilities. *Disability and Health Journal*, 8(2), 165-172.
- Lana-Elola, E., Watson-Scales, S. D., Fisher, E. M., & Tybulewicz, V. L. (2011). Down syndrome: Searching for the genetic culprits. *Disease Models & Mechanisms*, 4(5), 586-595.
- Landes, S. D., Stevens, J. D., & Turk, M. A. (2020). Cause of death in adults with Down syndrome in the United States. *Disability and Health Journal*, 13(4), 100947.
- Mandell, D. (2018). Dying before their time: Addressing premature mortality among autistic people. *Autism*, 22(3), 234-235.
- Mandell, D. S., Morales, K. H., Xie, M., Lawer, L. J., Stahmer, A. C., & Marcus, S. C. (2010). Age of diagnosis among Medicaid-enrolled children with autism, 2001-2004. *Psychiatric Services (Washington, D.C.)*, 61(8), 822-829.
- Medicaid and CHIP Payment and Access Commission (MACPAC). (2018). *People with disabilities*. <https://www.macpac.gov/subtopic/people-with-disabilities/>.
- Piven, J., Rabins, P., & Autism- in- Older Adults Working Group. (2011). Autism spectrum

- disorders in older adults: Toward defining a research agenda. *Journal of the American Geriatrics Society*, 59(11), 2151-2155.
- Ptomey, L. T., Walpitage, D. L., Mohseni, M., Dreyer Gillette, M. L., Davis, A. M., Forseth, B., ... & Waitman, L. R. (2020). Weight status and associated comorbidities in children and adults with Down syndrome, autism spectrum disorder and intellectual and developmental disabilities. *Journal of Intellectual Disability Research*, 64(9), 725-737.
- Ranjan, S., Nasser, J. A., & Fisher, K. (2018). Prevalence and potential factors associated with overweight and obesity status in adults with intellectual developmental disorders. *Journal of Applied Research in Intellectual Disabilities*, 31, 29-38.
- Rector, T. S., Wickstrom, S. L., Shah, M., Thomas Greenlee, N., Rheault, P., Rogowski, J., ... & Escarce, J. J. (2004). Specificity and sensitivity of claims- based algorithms for identifying members of Medicare+ Choice health plans that have chronic medical conditions. *Health Services Research*, 39(6p1), 1839-1858.
- Roux, A.M., Chvasta, K., Koffer M., Kaitlin H., Cooper, D., Tao, S., Assing-Murray, E., Shattuck, P., and Shea, L. (2023, March). National Autism Indicators Report: Introduction to Medicaid and Autism. Philadelphia, PA: Policy and Analytics Center, A.J. Drexel Autism Institute, Drexel University.
- Rubenstein, E., & Bishop, L. (2019). Is the autism boom headed for Medicaid? Patterns in the enrollment of autistic adults in Wisconsin Medicaid, 2008–2018. *Autism Research*, 12(10), 1541-1550.
- Schott, W., Tao, S., & Shea, L. (2022). Co- occurring conditions and racial- ethnic disparities: Medicaid enrolled adults on the autism spectrum. *Autism Research*, 15(1), 70-85.
- Snyder, L., & Rudowitz, R. (2015). Medicaid financing: How does it work and what are the

- implications. *Kaiser Family Foundation*. *kff.org/medicaid/issue-brief/medicaid-financing-how-does-itwork-and-what-are-the-implications/*. Retrieved August, 30, 2016.
- State Health Facts: Wisconsin: Medicaid & CHIP*. Kaiser Family Foundation (KFF). (n.d.). Retrieved March 3, 2023, from <https://www.kff.org/state-category/medicaid-chip/?state=WI>
- Status of state Medicaid Expansion Decisions: Interactive Map*. Kaiser Family Foundation (KFF). (2023, February 27). Retrieved March 3, 2023, from <https://www.kff.org/medicaid/issue-brief/status-of-state-medicaid-expansion-decisions-interactive-map/>
- Tye, C., Runicles, A. K., Whitehouse, A. J., & Alvares, G. A. (2019). Characterizing the interplay between autism spectrum disorder and comorbid medical conditions: An integrative review. *Frontiers in Psychiatry, 9*, 751.
- Tyler, C. V., Schramm, S. C., Karafa, M., Tang, A. S., & Jain, A. K. (2011). Chronic disease risks in young adults with autism spectrum disorder: forewarned is forearmed. *American Journal on Intellectual and Developmental Disabilities, 116*(5), 371-380.
- Vohra, R., Madhavan, S., & Sambamoorthi, U. (2017). Comorbidity prevalence, healthcare utilization, and expenditures of Medicaid enrolled adults with autism spectrum disorders. *Autism, 21*(8), 995-1009.
- Wang, L., & Leslie, D. L. (2010). Health care expenditures for children with autism spectrum disorders in Medicaid. *Journal of the American Academy of Child & Adolescent Psychiatry, 49*(11), 1165-1171.
- Weir, E., Allison, C., & Baron-Cohen, S. (2022). Autistic adults have poorer quality healthcare

and worse health based on self-report data. *Molecular Autism*, 13(1), 23.

<https://doi.org/10.1186/s13229-022-00501-w>

WHO, G. (2011). Global status report on noncommunicable diseases 2010.

Wisconsin Office of Rural Health. (2022, October 25). *WISH: Urban and rural counties*.

Wisconsin Department of Health Services. Retrieved February 28, 2023, from

<https://dhs.wisconsin.gov/wish/urban-rural.htm>


Zigman, W. B., Devenny, D. A., Krinsky- McHale, S. J., Jenkins, E. C., Urv, T. K., Wegiel, J.,


... & Silverman, W. (2008). Alzheimer's disease in adults with Down


syndrome. *International Review of Research in Mental Retardation*, 36, 103-145.


**Figures 1a-d**

*Wisconsin Medicaid beneficiaries (Aged 21+) with chronic health conditions, 2008-2018*










**Figure 2**

*Adjusted Odds for Selected Health Conditions in Adult Medicaid Beneficiaries Aged 21+ in Wisconsin*



**Table 1**

*Demographic characteristics of Wisconsin Medicaid beneficiaries (Aged 21+) with I/DD, 2008-2018*

|  | <b>Autism Only</b> | <b>Autism + Intellectual Disability</b> | <b>Down syndrome Only</b> | <b>Intellectual Disability Only</b> |
|--|--------------------|---|---------------------------|-------------------------------------|
|  | n=2828 (18.54%)    | n=1620 (10.62%)                         | n=2040 (13.37%)           | n=8769 (57.48%)                     |
| <b>Mean Age (SD)</b>   | 32.63 (11.72)      | 42.50 (15.12)                           | 46.79 (14.91)             | 52.21 (17.63)                       |
| <b>Age Groups</b>  |                    |   |                           |                                     |
| 21-34  | 2029 (71.75%)      | 653 (40.31%)                            | 558 (27.35%)              | 1763 (20.10%)                       |
| 35-49  | 522 (18.46%)       | 428 (26.42%)                            | 522 (25.59%)              | 2121 (24.19%)                       |
| 50-64  | 209 (7.39%)        | 391 (24.14%)                            | 713 (34.95%)              | 2714 (30.95%)                       |
| 65+  | 68 (2.40%)         | 148 (9.14%)                             | 247 (12.11%)              | 2171 (24.76%)                       |
| <b>Gender</b>  |                    |   |                           |                                     |
| Male   | 2064 (72.98%)      | 1122 (69.26%)                           | 1015 (49.75%)             | 4496 (51.27%)                       |
| Female   | 764 (27.02%)       | 498 (30.74%)                            | 1025 (50.25%)             | 4273 (48.73%)                       |
| <b>Race/Ethnicity</b>  |                    |   |                           |                                     |
| White (non-Hispanic)   | 1685 (59.58%)      | 1006 (62.10%)                           | 1359 (66.62%)             | 5491 (62.62%)                       |
| Black (non-Hispanic)   | 139 (4.92%)        | 122 (7.53%)                             | 63 (3.09%)                | 778 (8.87%)                         |
| Hispanic   | 68 (2.40%)         | 28 (1.73%)                              | 44 (2.16%)                | 211 (2.41%)                         |
| Asian or Pacific Islander; Or Native American or Alaskan Native (non-Hispanic) | 45 (1.59%)         | 25 (1.54%)                              | 28 (1.37%)                | 186 (2.12%)                         |
| Other or Unknown (non-Hispanic)  | 891 (31.51%)       | 439 (27.10%)                            | 546 (26.76%)              | 2099 (23.94%)                       |
| <b>County Designation</b>  |                    |   |                           |                                     |
| Urban  | 2052 (72.56%)      | 1197 (73.89%)                           | 1399 (68.58%)             | 6103 (69.60%)                       |
| Rural  | 776 (27.44%)       | 423 (26.11%)                            | 641 (31.42%)              | 2666 (30.40%)                       |

*Note.* Frequencies and percentages are reported for categorical variables. Means and standard deviations are reported for continuous variables. Rural and urban categories are based on the Wisconsin Office of Rural Health rural classifications.

**Tables 2a-2d**

*Wisconsin Medicaid beneficiaries (Aged 21+) with chronic health conditions, 2008-2018*

| <b>Autism Only</b>   |                 |                |                |             |                    |
|----------------------|-----------------|----------------|----------------|-------------|--------------------|
| <b>Age Groups</b>    | 21-34<br>n=2029 | 35-49<br>n=522 | 50-64<br>n=209 | 65+<br>n=68 | All Ages<br>n=2828 |
| <b>Asthma</b>        | 218 (10.74%)    | 86 (16.48%)    | 30 (14.35%)    | 15 (22.06%) | 349 (12.34%)       |
| <b>Diabetes</b>      | 237 (11.68%)    | 157 (30.08%)   | 108 (51.67%)   | 30 (41.12%) | 532 (18.81%)       |
| <b>Heart Disease</b> | 621 (30.61%)    | 275 (52.68%)   | 151 (72.25%)   | 56 (82.35%) | 1103 (39%)         |
| <b>Hypertension</b>  | 174 (8.58%)     | 131 (25.09%)   | 101 (48.33%)   | 38 (55.88%) | 444 (15.70%)       |

| <b>Autism + Intellectual Disability</b> |                |                |                |              |                    |
|---|----------------|----------------|----------------|--------------|--------------------|
| <b>Age Groups</b>                       | 21-34<br>n=653 | 35-49<br>n=428 | 50-64<br>n=391 | 65+<br>n=148 | All Ages<br>n=1620 |
| <b>Asthma</b>                           | 82 (12.56%)    | 50 (11.68%)    | 36 (9.21%)     | 17 (11.49%)  | 185 (11.42%)       |
| <b>Diabetes</b>                         | 127 (19.45%)   | 153 (35.75%)   | 140 (35.81%)   | 71 (47.97%)  | 491 (30.31%)       |
| <b>Heart Disease</b>                    | 299 (45.79%)   | 277 (64.72%)   | 307 (78.52%)   | 133 (89.86%) | 1016 (62.72%)      |
| <b>Hypertension</b>                     | 85 (13.02%)    | 124 (28.97%)   | 162 (41.43%)   | 79 (53.38%)  | 450 (27.78%)       |

| <b>Down Syndrome Only</b> |                |                |                |              |                    |
|---------------------------|----------------|----------------|----------------|--------------|--------------------|
| <b>Age Groups</b>         | 21-34<br>n=558 | 35-49<br>n=522 | 50-64<br>n=713 | 65+<br>n=247 | All Ages<br>n=2040 |
| <b>Asthma</b>             | 63 (11.29%)    | 77 (14.75%)    | 84 (11.78%)    | 35 (14.17%)  | 259 (12.70%)       |
| <b>Diabetes</b>           | 87 (15.59%)    | 160 (30.65%)   | 224 (31.42%)   | 79 (31.98%)  | 550 (26.96%)       |
| <b>Heart Disease</b>      | 237 (42.47%)   | 330 (63.22%)   | 592 (83.03%)   | 206 (83.40%) | 1365 (66.91%)      |
| <b>Hypertension</b>       | 43 (7.71%)     | 103 (19.73%)   | 218 (30.58%)   | 57 (23.08%)  | 421 (20.64%)       |

| <b>Intellectual Disability Only</b> |                 |                 |                 |               |                    |
|-------------------------------------|-----------------|-----------------|-----------------|---------------|--------------------|
| <b>Age Groups</b>                   | 21-34<br>n=1763 | 35-49<br>n=2121 | 50-64<br>n=2714 | 65+<br>n=2171 | All Ages<br>n=8769 |
| <b>Asthma</b>                       | 394 (22.35%)    | 478 (22.54%)    | 542 (19.98%)    | 357 (16.44%)  | 1171 (13.35%)      |
| <b>Diabetes</b>                     | 376 (21.33%)    | 804 (37.91%)    | 1261 (46.46%)   | 1097 (50.53%) | 3538 (40.35%)      |
| <b>Heart Disease</b>                | 967 (54.85%)    | 1510 (71.19%)   | 2175 (80.14%)   | 1950 (89.82%) | 6602 (75.29%)      |
| <b>Hypertension</b>                 | 262 (14.86%)    | 725 (34.18%)    | 1379 (50.81%)   | 1188 (54.72%) | 3554 (40.53%)      |

*Note.* These figures display the proportions of Wisconsin Medicaid beneficiaries (Aged 21+) with claims for asthma, diabetes, heart disease, and hypertension by age and diagnostic group, 2008-2018.

### Tables 3a-b

#### *Odds of Chronic Health Conditions for Adult Medicaid Beneficiaries, 2008-2018*

|  | Asthma            |                   | Diabetes          |                   |
|--|-------------------|-------------------|-------------------|-------------------|
|  | OR [95% CI]       | aOR [95% CI]      | OR [95% CI]       | aOR [95% CI]      |
| Disability Status (ref: autism only)   |                   |                   |                   |                   |
| Autism + Intellectual Disability   | 0.92 [.76-1.1]    | 0.92 [.76-1.1]    | 1.88 [1.6-2.2]*** | 1.41 [1.2-1.6]*** |
| Down syndrome  | 1.03 [.87-1.2]    | 0.99 [.83-1.2]    | 1.59 [1.4-1.8]*** | 1.06 [.92-1.2]    |
| Intellectual Disability Only   | 1.79 [1.6-2.0]*** | 1.74 [1.5-1.9]*** | 2.92 [2.6-3.2]*** | 1.65 [1.5-1.9]*** |
| Age  |                   | 0.99 [.99-.998]** |                   | 1.03 [1.0-1.1]*** |
| Gender (ref: male)   |                   |                   |                   |                   |
| Female   |                   | 1.55 [1.4-1.7]*** |                   | 1.09 [1.0-1.2]*   |
| Race/Ethnicity (ref: non-Hispanic White)                                       |                   |                   |                   |                   |
| Black (non-Hispanic)   |                   | 1.58 [1.4-1.8]*** |                   | 1.61 [1.4-1.8]*** |
| Hispanic   |                   | 0.92 [.69-1.2]    |                   | 1.54 [1.2-1.9]*** |
| Asian or Pacific Islander; Or Native American or Alaskan Native (non-Hispanic) |                   | 0.76 [.54-1.1]    |                   | 1.44 [1.2-1.9]**  |
| Other or Unknown (non-Hispanic)  |                   | 0.94 [.85-1.0]    |                   | 0.99 [.92-1.1]    |
| County Designation (ref: urban)  |                   |                   |                   |                   |
| Rural  |                   | 1.00 [.91-1.1]    |                   | 0.85 [.79-.92]*** |

|  | Heart Disease     |                   | Hypertension      |                     |
|--|-------------------|-------------------|-------------------|---------------------|
|  | OR [95% CI]       | aOR [95% CI]      | OR [95% CI]       | aOR [95% CI]        |
| Disability Status (ref: autism only)   |                   |                   |                   |                     |
| Autism + Intellectual Disability   | 2.63 [2.3-2.9]*** | 1.75 [1.5-1.9]*** | 2.07 [1.8-2.4]*** | 1.39 [1.2-1.6]***   |
| Down syndrome  | 3.16 [2.8-3.6]*** | 1.69 [1.5-1.9]*** | 1.39 [1.2-1.6]*** | 0.81 [.69-.94]**    |
| Intellectual Disability Only   | 4.76 [2.8-3.6]*** | 2.09 [1.9-2.3]*** | 3.66 [3.3-4.1]*** | 1.75 [1.5-1.9]***   |
| Age  |                   | 1.05 [1.0-1.1]*** |                   | 1.04 [1.03-1.04]*** |
| Gender (ref: male)   |                   |                   |                   |                     |
| Female   |                   | 1.19 [1.1-1.3]*** |                   | 0.91 [.84-.98]***   |
| Race/Ethnicity (ref: non-Hispanic White)                                       |                   |                   |                   |                     |
| Black (non-Hispanic)   |                   | 1.18 [1.0-1.4]*   |                   | 1.66 [1.4-1.9]***   |
| Hispanic   |                   | 0.98 [.77-1.2]    |                   | 1.21 [.94-1.6]      |
| Asian or Pacific Islander; Or Native American or Alaskan Native (non-Hispanic) |                   | 0.89 [.69-1.2]    |                   | 1.19 [.91-1.6]      |
| Other or Unknown (non-Hispanic)  |                   | 0.93 [.85-1.0]    |                   | 0.98 [.89-1.1]      |
| County Designation (ref: urban)  |                   |                   |                   |                     |
| Rural  |                   | 0.84 [.78-.92]*** |                   | 0.97 [.79-.94]**    |

*Note.* These tables display the unadjusted and adjusted odds ratios of asthma, diabetes, heart disease, and hypertension for Wisconsin Medicaid beneficiaries (21+) with IDD, 2008-2018. OR=unadjusted odds ratios. aOR= adjusted odds ratios, adjusting for age, race/ethnicity, gender, county of residence.

\* indicates  $p < .05$ , \*\* indicates  $p < .01$ , \*\*\* indicates  $p < .001$ .